

Anomalous Density-of-States Fluctuations in Two-Dimensional Clean Metals

T.R. Kirkpatrick¹ and D. Belitz²

¹*Institute for Physical Science and Technology and Department of Physics,
University of Maryland, College Park, MD 20742*

²*Department of Physics and Theoretical Science Institute, University of Oregon, Eugene, OR 97403*

(Dated: August 22, 2012)

It is shown that density-of-states fluctuations, which can be interpreted as the order-parameter susceptibility χ_{OP} in a Fermi liquid, are anomalously strong as a result of the existence of Goldstone modes and associated strong fluctuations. In a 2- d system with a long-range Coulomb interaction, a suitably defined χ_{OP} diverges as $1/T^2$ as a function of temperature in the limit of small wavenumber and frequency. In contrast, standard statistics suggest $\chi_{\text{OP}} = O(T)$, a discrepancy of three powers of T . The reasons behind this surprising prediction, as well as ways to observe it, are discussed.

PACS numbers:

It is known that the Fermi-liquid (FL) state can be described in terms of a stable renormalization-group (RG) fixed point (FP) [1]. A related concept is the possibility of understanding the FL as an ordered state, in analogy to the ferromagnetically ordered state and its corresponding stable RG FP [2, 3]. This notion was pioneered by Wegner in the context of disordered systems [4]. More recently it was realized that the principle behind it is much more general and powerful, and can be applied to clean FLs as well [5, 6]. Important features of the resulting effective field theory are that it integrates out all massive degrees of freedom to arrive at an effective soft-mode theory that allows for a very straightforward analysis of the FL FP, associated corrections to scaling, and the scaling behavior of various observables. Within this conceptual framework, the order parameter (OP) is the density of states (DOS) [7], the field conjugate to the OP is a frequency-dependent chemical potential, the spontaneously broken continuous symmetry is a rotational symmetry in frequency space that can be understood as the symmetry between retarded and advanced degrees of freedom, and the resulting Goldstone modes are the soft particle-hole excitations. These quantities are analogous to the magnetization m , the magnetic field h , the spin-rotational symmetry, and the magnons, respectively, in a Heisenberg ferromagnet. Note, however, that the Goldstone modes in the FL case are soft only at zero temperature ($T = 0$).

Let us recall the behavior of the DOS as a function of the frequency ω (or the energy distance from the Fermi surface). For our purposes we mainly focus on two-dimensional (2- d) systems with a long-ranged Coulomb interaction, which is arguably the most interesting case. However, the theory can be generalized to arbitrary $d > 1$, and the case of a short-ranged interaction is also of interest, see below and Ref. 8. It is known that the DOS is a nonanalytic function of ω , namely,

$$N(\omega) = 2N_{\text{F}}(1 + a|\omega|/\epsilon_{\text{F}}) + o(\omega) \quad (1)$$

Here N_{F} is the DOS per spin at the Fermi level ϵ_{F} , and

a is a dimensionless coefficient. $o(\omega)$ denotes terms that vanish faster than linearly for $\omega \rightarrow 0$. This result follows from the soft-mode theory developed in Refs. 5, 6 (see below); technically, the frequency correction in Eq. (1) is the leading correction to scaling at the FL FP. It has been calculated explicitly within many-body perturbation theory, which yields a weak-coupling value for a , viz., $a = 1/4$ [9–12]. It is important to note, however, that perturbation theory alone can never establish the $|\omega|$ as the true leading frequency dependence. Our RG analysis of the soft-mode field theory, on the other hand, guarantees that the exponent of unity is exact. The prefactor a will acquire corrections if one goes to higher-loop order. The linear frequency dependence is also consistent with simple scaling considerations [6], as it should be [13]. It has been observed in high-mobility tunnel junctions [14].

With these results as a backdrop, it is very interesting to ask about the behavior of the OP susceptibility. This quantity is an important measure of the strength of fluctuations in the ordered phase, which in turn provides a measure of the likelihood that the ordered phase will become unstable as a function of a suitable parameter. In the case of the classical Heisenberg ferromagnet mentioned above, the OP susceptibility is the magnetic susceptibility $\chi_{\text{m}} = \partial m / \partial h$. As a function of the wave number k , the longitudinal component of χ_{m} diverges for $k \rightarrow 0$ for all dimensions $d < 4$: $\chi_{\text{m,L}}(k) \propto 1/k^{4-d}$ [15]. This reflects the strong fluctuations in the ordered phase due to the existence of Goldstone modes that couple to the OP. It is then natural to ask whether an analogous susceptibility exists in the FL, and what its behavior at small wave numbers and frequencies is.

To answer this question, we have performed explicit calculations [8], some of which are quoted below. However, for the sake of physical clarity it is very illustrative to derive the results from qualitative arguments, which is the purpose of this Letter. To this end, we first represent the DOS as

$$N(\omega) = \text{Re } G(\mathbf{x} = 0, i\omega_n \rightarrow \omega + i0)/\pi. \quad (2)$$

Here

$$G(\mathbf{x} = 0, i\omega_n) = \langle \rho(\mathbf{x}, i\omega_n) \rangle \quad (3a)$$

is the exact Green's function, with

$$\rho(\mathbf{x}, i\omega_n) = \bar{\psi}_n(\mathbf{x}) \psi_n(\mathbf{x}) \quad (3b)$$

in terms of fermion fields $\bar{\psi}_n(\mathbf{x})$ and $\psi_n(\mathbf{x})$ that depend on a fermionic Matsubara frequency $\omega_n = 2\pi T(n + 1/2)$, ($n = 0, \pm 1, \dots$). We suppress the spin degree of freedom, which is not important for our purposes. $\rho(\mathbf{x}, i\omega_n)$ can be interpreted as the density of particles at point $\mathbf{x}$ and (complex) energy $i\omega_n$ [16]. The temporal Fourier transform is defined as

$$\psi_n(\mathbf{x}) = T^{1/2} \int_0^{1/T} d\tau e^{i\omega_n \tau} \psi(\mathbf{x}, \tau), \quad (4)$$

with τ the imaginary time variable. Dimensionally, $\psi(\mathbf{x}, \tau)$ is an inverse square root of a volume, so N is dimensionally indeed a DOS. Now we consider the spatial Fourier transform $\rho(\mathbf{k}, i\omega_n) = \int d\mathbf{x} e^{-i\mathbf{k} \cdot \mathbf{x}} \rho(\mathbf{x}, i\omega_n)$, and the fluctuation $\delta\rho = \rho - \langle \rho \rangle$, and define a correlation function

$$\chi_{\text{OP}}(\mathbf{k}, i\omega_n) = \langle \delta\rho(\mathbf{k}, i\omega_n) \delta\rho(-\mathbf{k}, i\omega_n) \rangle. \quad (5)$$

χ_{OP} is the order-parameter susceptibility of interest. It describes both the response of the OP (i.e., the DOS) to its conjugate field, and the spontaneous fluctuations of the OP. The former is related to the latter by the fluctuation-dissipation theorem.

Let us first use simple statistical arguments to determine the behavior of χ_{OP} one would expect in the absence of anomalous fluctuations. This will also serve to make a general point that is related to the definition of the temporal Fourier transform above. Consider $\varphi_n(\mathbf{x}) = \psi_n(\mathbf{x})/\sqrt{T}$, and define the “volume” $V_T \equiv 1/T$ in the imaginary-time direction of the space-time of quantum statistical mechanics. Then $\langle p_n(\mathbf{x}) \rangle \equiv \langle \bar{\varphi}_n(\mathbf{x}) \varphi_n(\mathbf{x}) \rangle \propto V_T$ is a “time-extensive” quantity that is proportional to $V_T \equiv 1/T$. Now consider the relative fluctuation $\langle (\delta p_n(\mathbf{x}))^2 \rangle / \langle p_n(\mathbf{x}) \rangle^2 \propto V_T / V_T^2 = 1/V_T = T$. This just says that the relative fluctuation is proportional to $1/V_T$, as one would expect from ordinary statistics. This yields an estimate for the fluctuations of ρ :

$$\langle (\delta\rho_n(\mathbf{x}, i\omega_n))^2 \rangle \propto \frac{\langle (\delta p_n)^2 \rangle}{V_T^2} \propto \frac{\langle (\delta p_n)^2 \rangle / V_T^2}{(\langle p_n \rangle / V_T)^2} \propto 1/V_T = T. \quad (6)$$

All of these simple arguments of course assume that there are no strong fluctuations in the system that invalidate the simple statistics. Independent of this assumption, however, they show that $\langle (\delta\rho_n(\mathbf{x}, i\omega_n))^2 \rangle$, and hence χ_{OP} , must carry a factor of T that is simply the inverse linear extension $1/V_T$ of the imaginary-time axis. We thus have

$$\chi_{\text{OP}}(\mathbf{k}, i\omega_n; T) = T \theta(\mathbf{k}, i\omega_n; T). \quad (7)$$

In the absence of anomalous fluctuations, θ will scale as the zeroth power of the wave number, the frequency, or the temperature; i.e., $\theta \sim 1$.

It is easy to see that the latter conclusion cannot be correct in an actual FL. Consider the scaling arguments from Ref. 17, and first consider the case of a short-ranged interaction, in which case there is only one dynamical scaling exponent $z = 1$. This reflects the fact that in the Goldstone modes that characterize the broken symmetry in the FL, the frequency scales as the wave number, $\omega \sim k$. At the stable FL FP, and in $d = 2$, the scaling part of the free energy density f obeys the homogeneity law [18]

$$f(T, h) = b^{-(2+z)} f(Tb^z, hb^z). \quad (8)$$

Here we have assigned scale dimensions $[L] = -1$ and $[T] = z$ to factors of length and temperature (or energy, frequency, or inverse time), respectively. b is the RG length rescaling factor. h is the field conjugate to the OP, which has a scale dimension $[h] = z$ since the DOS must scale as an inverse energy times an inverse volume. Frequency-dependent quantities will scale with respect to frequency the same as with respect to temperature. Differentiating once with respect to h , we obtain for the scaling part of the OP density, i.e., the DOS correction $\delta N = N - 2N_F$,

$$\delta N(T) = b^{-1} \delta N(Tb) \propto T, \quad (9)$$

in agreement with Eq. (1). Differentiating again with respect to h yields

$$\chi_{\text{OP}}(T, k) = \chi_{\text{OP}}(Tb, kb) = f_\chi(T/k), \quad (10)$$

with f_χ a scaling function. If we combine this with the conclusion from Eq. (7), namely, that χ_{OP} must be proportional to T for fundamental quantum statistical reasons, we conclude

$$\chi_{\text{OP}}(T, k) \propto \begin{cases} T/k & \text{for } T \rightarrow 0 \text{ at fixed } k \\ T/T = O(1) & \text{for } k \rightarrow 0 \text{ at fixed } T. \end{cases} \quad (11)$$

This result is not consistent with the naive statistical arguments given above: If the DOS were normally distributed, we would have $\chi_{\text{OP}} \propto 1/V_T$ and $\theta \sim 1$. Instead, we see that the quantity θ in Eq. (7) scales as $\theta \sim 1/k \sim 1/T$, and $\chi_{\text{OP}} \sim 1$. This divergence of the relative DOS fluctuations reflects the strong fluctuations in the system that are a consequence of the existence of the Goldstone modes.

In order to understand what happens in the case of a long-ranged Coulomb interaction, we consider the structure of the field theory developed in Refs. 5, 6. Within this framework, the DOS is schematically given by a frequency-momentum integral over a vertex function $V(k, i\omega)$ times the potential $U(k, i\omega)$, with k the wave number and $i\omega$ the imaginary frequency. The vertex

function has the structure $V(k, i\omega) = v(\omega/v_F k)/k^2$, with $v(x \rightarrow \infty) \propto 1/x^2$. U is a constant in the short-range case, and represents the dynamically screened Coulomb potential in the long-range case. In $d = 2$, the latter has the schematic structure

$$U(k, i\omega) = \frac{1}{k + \kappa - \frac{\kappa\omega/v_F k}{\sqrt{1+\omega^2/(v_F k)^2}}} \approx \frac{1}{k + \kappa(v_F k)^2/\omega^2}. \quad (12)$$

Here κ is the screening wave number, and the second expression is valid in the limit $v_F k \ll \omega$. In the denominator we recognize the plasmon excitation, with its characteristic $\omega \sim k^{1/2}$ scaling. We also see the well-known fact that screening breaks down at nonzero frequencies.

The DOS is schematically given by an integral [19]

$$N \sim \int_0^\Lambda dk k \int_T^{v_F \Lambda} d\omega \frac{1}{k^2} v(\omega/v_F k) U(k, i\omega), \quad (13)$$

with Λ an ultraviolet momentum cutoff. Power counting yields $N(T) \propto \text{const.} - T$, in agreement with Eqs. (9) and (1). This is true in both the short-range and the long-range case; the singular nature of the screened Coulomb interaction at nonzero frequency does not suffice to change the behavior in $d = 2$ [20].

Now consider the OP susceptibility. It follows from Eq. (5) that χ_{OP} is given by an integral analogous to the one in Eq. (13) with the integrand squared and an extra factor of T :

$$\chi_{\text{OP}} \sim T \int_0^\Lambda dk k \int_T^{v_F \Lambda} d\omega \left(\frac{1}{k^2} v(\omega/v_F k) U(k, i\omega) \right)^2. \quad (14)$$

The prefactor of T is just the trivial “volume” factor $1/V_T$ discussed in the context of Eqs. (6, 7). In the short-range case, we obtain $\chi_{\text{OP}} \propto T/T$ in agreement with Eq. (11). In the long-range case, a power-counting analysis of the integral yields $\chi_{\text{OP}} \propto T/T^3$, with a multiplicative logarithmic infinity coming from the momentum integral.

A careful analysis of χ_{OP} within the field theory of Ref. 6, or its generalization to a long-ranged Coulomb interaction, yields results that are consistent with the ones sketched above. For the static OP susceptibility in the limit of low temperatures we find, in the long-range case,

$$\chi_{\text{OP}}(k) = \frac{\kappa^2}{90\pi} \frac{T}{T^3} \ln(T^2/v_F^2 \kappa k) \quad (15a)$$

This result is valid for $T^2/v_F k \gg v_F \kappa$, and to logarithmic accuracy. In the opposite limit, we find

$$\chi_{\text{OP}}(k) = \text{const.} \times \frac{\kappa^{1/2}}{v_F^3} \frac{T}{k^{3/2}}, \quad (15b)$$

with const. a constant of $O(1)$ [21]. Finally, for $T \rightarrow 0$ and nonzero external frequency ω one finds, to logarithmic accuracy for $\omega^2 \gg v_F^2 \kappa k$,

$$\chi_{\text{OP}}(k, i\omega) = \frac{8}{3\pi^2} \frac{\kappa^2 T}{\omega^3} \ln(\omega^2/v_F^2 \kappa k) \quad (15c)$$

As expected, the frequency scaling is the same as the temperature scaling. Comparing Eqs. (15b) and (15a, 15c) we see that T or ω scale as $k^{1/2}$, which reflects the fact that the integral is dominated by the plasmon time scale. This is also apparent from the arguments of the logarithms in Eqs. (15a, 15c). The fact that $\chi_{\text{OP}}(k \rightarrow 0)$ diverges even at a nonzero T is due to the fact that the plasmon, being a density fluctuation governed by a conservation law, remains soft even at $T > 0$. Consequently, there is a logarithmic divergence (in $d = 2$) that is not cut off by temperature.

Also of interest is the homogeneous susceptibility in a finite system of linear dimension L . In that case, k in Eqs. (15) gets replaced by $1/L$, and we have

$$\chi_{\text{OP}} \propto \begin{cases} (1/T^2) \ln(T^2 L/v_F^2 \kappa) & \text{for } T^2 L \gg v_F^2 \kappa \\ T L^{3/2} \kappa^{1/2}/v_F^3 & \text{for } T^2 L \ll v_F^2 \kappa \\ (T/\omega^3) \ln(\omega^2 L/v_F^2 \kappa) & \text{for } T \rightarrow 0, \omega^2 L \gg v_F^2 \kappa \end{cases}. \quad (16)$$

In summary, we have calculated the susceptibility that describes density-of-states fluctuations in a 2 - d Fermi liquid; a correlation function that can be interpreted as the OP susceptibility of the ordered phase represented by the Fermi liquid. The OP susceptibility, properly normalized to account for a trivial factor of T , see Eq. (7), diverges in the limit of vanishing wave number k as $1/k^{3/2}$, Eq. (15b), or as $1/T^3$ or $1/\omega^3$ in the limits of vanishing temperature or frequency, respectively. This is a very strong effect; uncorrelated statistics would lead to a constant for the same quantity. It is the combined result of the Goldstone modes, i.e., particle-hole excitations, in the ordered phase, and the plasmon excitations in an electron system with long-range Coulomb interaction. The frequency scale associated with the latter ($\omega \sim k^{1/2}$), although formally irrelevant compared to the Goldstone mode frequency scale ($\omega \sim k$), acts as a dangerous irrelevant variable in the RG sense and leads to an even stronger divergence than one would expect from naive scaling. Even in the case of a short-range interaction [22], where only the Goldstone modes are present and naive scaling works, the normalized susceptibility diverges as $1/k \sim 1/T \sim 1/\omega$, compared to a constant in the absence of anomalous fluctuations. A simple analogy of this effect in a classical system is the $1/k^{4-d}$ divergence of the longitudinal spin susceptibility in the ordered phase of a Heisenberg ferromagnet.

Physically, this anomaly is very closely related to the $|\omega|$ anomaly in the DOS, Eq. (1), which constitutes a pseudogap as originally defined by Mott [23]. It is worth noting that this pseudogap is an intrinsic feature of *any* 2 - d Fermi liquid. We stress again that the current theory shows the linear frequency dependence to be exact, which cannot be established within perturbation theory.

We finally add some comments about the experimental relevance of the above results. The spectrum of the local Green's function, Eq. (3a), is what is often referred to as

the local density of states (LDOS) [24, 25]. The LDOS gives the dominant contribution to the tunneling current in a scanning tunneling experiment [26]; its spatial average, which is the DOS, is measured in a classic tunnel junction. Our OP susceptibility, Eq. (5), describes the spatially averaged fluctuations of the LDOS. A two-tip tunneling experiment [24, 27] should be able to give information about this quantity [28].

We are grateful to S. Gregory and H. Manoharan for helpful discussions. This work was supported by the National Science Foundation under Grant Nos. DMR-09-29966, DMR-09-01907, and PHY-10-66293. Part of this work was performed at the Aspen Center for Physics.

-
- [1] R. Shankar, *Rev. Mod. Phys.* **66**, 129 (1994).
[2] K. G. Wilson and J. Kogut, *Phys. Rep.* **12**, 75 (1974).
[3] S.-K. Ma, *Modern Theory of Critical Phenomena* (Benjamin, Reading, MA, 1976).
[4] F. Wegner, *Z. Phys. B* **35**, 207 (1979).
[5] D. Belitz and T. R. Kirkpatrick, *Phys. Rev. B* **56**, 6513 (1997).
[6] D. Belitz and T. R. Kirkpatrick, *Phys. Rev. B* **85**, 125126 (2012).
[7] More precisely, the single-particle spectral density, whose zeroth moment with respect to the momentum is the DOS, is the OP density. For simplicity, we will refer to the DOS as the OP.
[8] D. Belitz and T.R. Kirkpatrick, unpublished results.
[9] D. V. Khveshchenko and M. Reizer, *Phys. Rev. B* **57**, R4245 (1998).
[10] E. G. Mishchenko and A. V. Andreev, *Phys. Rev. B* **65**, 235310 (2002).
[11] Reference 9 actually obtained $a = 1/2$. To us, this seems to be due to a simple algebraic mistake, rather than a missing physical contribution as suggested in Ref. 10.
[12] This result holds in the long-ranged case. For remarks regarding the short-ranged case, see Ref. 20.
[13] In generic dimensions, scaling yields $\delta N \equiv N(\omega) - 2N_F \propto |\omega|^{d-1}$ [5, 6]. This agrees with an explicit calculation for all $d > 1$ in the case of a short-ranged interaction. For a Coulomb interaction, it agrees for all $d \geq 2$, but not for $1 < d < 2$, where the additional time scale provided by the plasmon serves as a dangerous irrelevant variable and leads to a stronger dependence, $\delta N \propto |\omega|^{(d-1)/(3-d)}$. Also, scaling is not sensitive to logarithmic corrections; e.g., in $d = 3$ one finds $\delta N \propto \omega^2 \log |\omega|$ [8].
[14] See Ref. 29. The interpretation given in this paper was in terms of different physics, in our opinion erroneously so. We also note that the pseudogap in the normal phase of the cuprates has recently been found to not scale with the superconducting properties, Ref. 30. The energy scale in the latter experiment is roughly consistent with Eq. (1), as is the one in Ref. 29. This raises the possibility that the pseudogap in the cuprates can also be explained in terms of 2- d Fermi-liquid physics. This notion requires further investigation.
[15] E. Brézin and D. J. Wallace, *Phys. Rev. B* **7**, 1967 (1973).
[16] Note, however, that ρ is defined as a product of two fermion fields at a frequency ω_n , whereas the actual particle number density would involve a convolution in frequency space.
[17] T. R. Kirkpatrick and D. Belitz, *Phys. Rev. Lett.* **108**, 086404 (2012).
[18] Here we just make a phenomenological scaling assumption. However, the homogeneity law, Eq. (8), and all of its consequences can be derived from the effective field theory, Refs. 5, 6, 8. In particular, this derivation makes it clear that the linear dependences in Eqs. (1) and (9) reflect the leading correction to scaling at the FL FP.
[19] This, as well as the structure of the vertex, can be seen from Eq. (5.4) in Ref. 6.
[20] For $d < 2$, the long-range potential changes the leading temperature or frequency dependence from T^{d-1} to $T^{(d-1)/(3-d)}$, as can also easily be seen by power counting. In a RG language, the plasmon time scale acts as a dangerous irrelevant variable that invalidates naive scaling. We also note that in $d = 2$ in the short-range case, a one-loop calculation within the field theory, as opposed to the power-counting arguments presented here, shows that the T or $|\omega|$ term has a zero prefactor [6], in agreement with earlier results from many-body perturbation theory [10, 31]. A complete one-loop calculation that takes into account the particle-particle channel shows that the leading contribution is proportional to $|\omega|/\ln \omega|^3$ [8]. The question whether at higher-loop order there is a nonvanishing pure $|\omega|$ contribution (with no logarithmic corrections) is an interesting open problem.
[21] In this case the prefactor is difficult to determine, since the wave number integral involves a convolution.
[22] A 2- d electron system with a short-range interaction can be realized, e.g., by gating a field-effect transistor, see Ref. 32.
[23] N. F. Mott, *Rev. Mod. Phys.* **40**, 677 (1968).
[24] Y. S. Chan and E. J. Heller, *Phys. Rev. Lett.* **78**, 2570 (1997).
[25] This definition of the LDOS is more general than the one found in earlier papers, e.g., Ref. 26, as it is in terms of the exact Green's function and thus includes the full many-body effects. We also note that our local Green's function is independent of the position because we consider a homogeneous model; it becomes position dependent if one takes into account the underlying lattice.
[26] J. Tersoff and D. R. Hamann, *Phys. Rev. B* **31**, 805 (1985).
[27] Q. Niu, M. C. Chang, and C. K. Shih, *Phys. Rev. B* **51**, 5502 (1995).
[28] No detailed theory of two-tip tunneling seems to exist. However, since tunneling is clearly sensitive to the density of electrons at a given energy, as evidenced by the sensitivity of one-tip tunneling to the LDOS, or the average of ρ as defined in Eqs. (3), it is plausible that a two-tip experiments will be sensitive to the fluctuations of ρ .
[29] P. Jiang, I. Yang, W. Kang, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, *Phys. Rev. Lett.* **96**, 126804 (2006).
[30] T. Jacobs, S. O. Katterwe, H. Motzkau, A. Rydh, A. Maljuk, T. Helm, C. Putzke, E. Kampert, M. V. Kartsovnik, and V. M. Krasnow (2012), arXiv:1202.6370.
[31] A. Chubukov, D. Maslov, S. Gangadharaiah, and L. Glazman, *Phys. Rev. B* **71**, 205112 (2005).
[32] T. Ando, A. B. Fowler, and F. Stern, *Rev. Mod. Phys.* **54**, 437 (1982).